

Testosterone gel (Tostran 2%) for low sex drive in menopausal women

Gynaecology Department

Information for Patients

Produced: January 2020

Review: January 2023

Leaflet number: 878 Version: 1

Introduction

This leaflet provides information about use of testosterone gel **Tostran 2%**, for menopausal women who have a low sex drive (libido). Testosterone is currently not licensed to women in the UK. The use of this gel is 'off label' which means that the manufacturer of the medicine has not applied for a license or has not specified that it can be used in this way. However, it has proven benefits in numerous clinical trials and is also recommended by the British Menopause Society.

What is testosterone?

Testosterone is one of the sex hormones produced naturally from ovaries and adrenal glands in women. It is commonly thought of as a male hormone but testosterone plays an important role in women's life including desire for sex. Around 50% of testosterone is produced by the ovaries and 50% by the adrenal glands. In fact, women produce 3 times more testosterone than oestrogen before the menopause.

Levels of testosterone gradually go down as women get older, but there is a sudden loss of testosterone if you have an operation to remove your ovaries (oophorectomy).

What are the potential benefits of testosterone gel?

Testosterone may improve libido, mood, energy and concentration. Low libido may improve with oestrogen therapy alone but in some cases, testosterone may be more helpful, especially in women who have had their ovaries removed. According to the National Institute for Health and Care Excellence (NICE) guidelines on menopause management, use of testosterone for women who are already taking hormone replacement therapy (HRT) may be helpful. It can sometimes take a few months for the full effects of testosterone to work.

**Health information and support is available at www.nhs.uk
or call 111 for non-emergency medical advice**

Visit www.leicestershospitals.nhs.uk for maps and information about visiting Leicester's Hospitals
To give feedback about this information sheet, contact InformationForPatients@uhl-tr.nhs.uk

How is testosterone gel used?

Tostran 2% testosterone gel comes in a canister. Pressing the pump once releases a pea sized amount equal to a 10mg dose which needs to be applied **every other day**.

- The gel should be rubbed onto dry, non-hairy areas such as the inner surface of forearms or thighs.
- The area should be allowed to dry before getting dressed and should not be washed for 2 to 3 hours after applying the gel.
- You should keep changing the area where you apply the gel to avoid hair growth in that area.
- Hands should be washed immediately after applying the gel to the body.
- The area of your skin should not come into contact with partners, children and pregnant women.

Do not use the amounts stated in the packet's leaflet as this is how much a man should use.

What are the side effects?

Clinical studies have shown that if you follow the instructions carefully, most people can use testosterone gel. Possible side effects are:

- increased facial or body hair (hirsutism) - uncommon
- male pattern hair loss (alopecia) which could be patchy or at the front - uncommon
- acne and greasy skin - uncommon
- deepening of voice - rare
- enlarged clitoris - rare

So far clinical studies have not shown an increased risk of heart disease or breast cancer, but more research is required in this field.

What blood test monitoring will be needed?

Blood testing for testosterone is not essential but can be useful.

Professionals have not yet agreed on what the 'normal' level of testosterone should be for women.

The Free Androgen Index (FAI) measurement can be useful to see whether your testosterone levels have gone up, however monitoring changes in your symptoms and side effects may be a better measure of success. If you find that testosterone is helpful then you would normally continue to use this while you are taking the standard HRT.

When should testosterone gel not be used?

Testosterone should be avoided:

- during pregnancy or breastfeeding
- if you have active liver disease
- if you have a history of hormone sensitive breast cancer
- if you are a competitive athlete
- for women with upper normal limits or high baseline free testosterone levels / FAI (>5%)

Contact details

You can contact the secretary of your menopause specialist at Leicester Royal Infirmary on 0116 258 6441.

Further information

The British Menopause Society website:

<https://thebms.org.uk/publications/tools-for-clinicians/testosterone-replacement-in-menopause/>

اگر آپ کو یہ معلومات کسی اور زبان میں درکار ہیں، تو براہ کرم مندرجہ ذیل نمبر پر ٹیلی فون کریں۔
على هذه المعلومات بلغةٍ أخرى، الرجاء الاتصال على رقم الهاتف الذي يظهر في الأسفل

જો તમને અન્ય ભાષામાં આ માહિતી જોઈતી હોય, તો નીચે આપેલ નંબર પર કૃપા કરી ટેલિફોન કરો

ਜੇ ਤੁਸੀਂ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਭਾਸ਼ਾ ਵਿਚ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਹੇਠਾਂ ਦਿੱਤੇ ਗਏ ਨੰਬਰ 'ਤੇ ਟੈਲੀਫੋਨ ਕਰੋ।

Aby uzyskać informacje w innym języku, proszę zadzwonić pod podany niżej numer telefonu

If you would like this information in another language or format such as EasyRead or Braille, please telephone 0116 250 2959 or email equality@uhl-tr.nhs.uk